

A new criterion for finite non-cyclic groups ^{*}

Wei Zhou

College of Mathematics & Finance, Southwest China Normal University , Chongqing 400715

People's Republic of China

E-mail: zh_great@hotmail.com

and

Wujie Shi

School of Mathematics, Suzhou University, Suzhou 215006

People's Republic of China

E-mail: wjshi@suda.edu.cn

and

Zeyong Duan

College of Mathematics & Finance, Southwest China Normal University , Chongqing 400715

People's Republic of China

Abstract

Let H be a subgroup of a group G . We say that H satisfies the power condition with respect to G , or H is a power subgroup of G , if there exists a non-negative integer m such that $H = G^m = \langle g^m | g \in G \rangle$. In this note, the following theorem is proved: Let G be a group and k the number of non-power subgroups of G . Then (1) $k = 0$ if and only if G is a cyclic group (theorem of F. Szász); (2) $0 < k < \infty$ if and only if G is a finite non-cyclic group; (3) $k = \infty$ if and only if G is a infinite non-cyclic group. Thus we get a new criterion for the finite non-cyclic groups.

Keywords: power subgroup, cyclic group, Dedekind group.

2000MR subject classification: 20E07, 20E34, 20D25

1. Introduction

Let H be a subgroup of group G . We say that H satisfies the power condition with respect to G , or H is a power subgroup of G if there exists a non-negative integer m such

^{*}Project supported by the National Natural Science Foundation(Grant No.10171074).

that $H = G^m = \langle g^m \mid g \in G \rangle$. On the other hand, if $H \neq G^m$ for all m , we say that H is a non-power subgroup of G . If H is a power subgroup of G , the power exponent is the least non-negative integer m such that $H = G^m$.

It is obvious that a nontrivial group G has at least two trivial power subgroups: $\{1\}$ and G itself, and the power exponents of the two subgroups are 0 and 1, respectively.

The power subgroups have some properties. For example, if H is a power subgroup of group G , then H is a full-invariant subgroup, in particular H is normal in G ; and if H is a non-power subgroup of G , then a conjugate subgroup $H^x (x \in G)$ is also a non-power subgroup of G .

The number of non-trivial power subgroups affects the structure of the group. In [2], F.Szász proved that G is a cyclic group if and only if all subgroups of G are power subgroups. That is, a non-cyclic group contains at least one non-power subgroup. In this paper, we generalize the above result of F. Szász and prove the following theorem:

Theorem. *Let G be a group and k the number of non-power subgroups of G . Then*

- (1) $k = 0$ if and only if G is a cyclic group;
- (2) $0 < k < \infty$ if and only if G is a finite non-cyclic group;
- (3) $k = \infty$ if and only if G is a infinite non-cyclic group.

Remark 1. *The conclusion (1) of this theorem is the theorem of F.Szász in [2].*

Remark 2. *Note that for some k in the theorem, the group G may not exist. In [3],[4] the author proved that the case of $k = 1$ or $k = 2$ do not occur. But there are groups having exactly 3 non-power subgroups: For example the quaternion group Q_8 of order 8, and $Z_2 \times Z_2$ are groups possessing just 3 non-power subgroups.*

From the theorem the following corollary gives a new criterion for finite non-cyclic groups.

Corollary. *Suppose G is a non-cyclic group. Then G is finite if and only if G contains only finitely many non-power subgroups.*

Problem. *For any integer $k (k \geq 3)$, does there exist groups possessing just k non-power subgroups?*

In the proof of the Theorem, we use the structure of the Dedekind group. A group is called Dedekind group if all its subgroups are normal. From Theorem 5.3.7 in [1], we know that G is a Dedekind group if and only if G is abelian or the direct product of a quaternion group of order 8, an elementary abelian 2-group and an abelian group with all its elements of odd order. Notation is standard and may be found for instance in [1]. In particular, we denote k the number of non-power subgroups of G .

2. Some lemmas

Lemma 1. *Let A be an abelian power subgroup in a group G . Then the set of power subgroups of G contained in A coincides with the set of power subgroups of A .*

Proof: If $G = A$, then there it is nothing to prove. Let $G \neq A$ and let l be the power exponent of A . Then $l > 1$ and $A = G^l$. Suppose that $G^m \leq A$ for $m \geq 1$ and let d be the greatest common divisor of l, m . Then $d = lr + ms$ for some integers r and s , and $g^d = (g^l)^r (g^m)^s \in A$ for every $g \in G$, so $G^d \leq A$. It is obvious that $A = G^l \leq G^d$, so $d = l$ and l divides m . Hence $m = ls_0$ for some $s_0 \geq 0$. Note that $A = G^l = \langle g^l | g \in G \rangle$ is abelian. We get $A^{s_0} = (G^l)^{s_0} = \langle g^l | g \in G \rangle^{s_0} = \langle g^{ls_0} | g \in G \rangle = G^m$. On the other hand, if A^n is a power subgroup of A , then $A^n = G^{ln}$ is a power subgroup of G . The lemma is proved.

Lemma 2. *If A/N is a non-power subgroup of a factor-group G/N , then A is a non-power subgroup of G .*

Proof: Suppose the result is false. That is, A is a power subgroup of G . Let $A = G^m = \langle g_i^m | g_i \in G, i = 1, 2, \dots \rangle$. Since $g_1^m \cdots g_r^m N = (g_1 N)^m \cdots (g_r N)^m$, $A/N = (G/N)^m$, a contradiction. Hence the lemma is proved.

Now we need a lemma about the structure of the cyclic group.

Lemma 3. *If G is an infinite abelian group all of whose proper quotient groups are finite, then G is infinite cyclic.*

Proof: Suppose G is not infinite cyclic. We choose an element $a_0 \in G$ and $a_0 \neq 1$. $\langle a_0 \rangle$ is a proper subgroup of G and $\langle a_0 \rangle \neq 1$. Then $G/\langle a_0 \rangle$ is finite. So $G/\langle a_0 \rangle$ is finitely generated and G is finitely generated. Then G is a direct product of finitely many cyclic groups of infinite or prime-power orders. Let $G = \langle g_1 \rangle \times \langle g_2 \rangle \cdots \times \langle g_n \rangle$, and $n > 1$ by the assumption. If there exists an i , such that $|g_i| = \infty$. Then $G/\langle g_1 \rangle \times \cdots \times \langle g_{i-1} \rangle \times \langle g_{i+1} \rangle \times \cdots \times \langle g_n \rangle \cong \langle g_i \rangle$ is finite by the condition. That is impossible. So $|g_i| < \infty, \forall i$, which makes G be a finite group, a contradiction. We prove the lemma.

Lemma 4. *If the number of cyclic subgroups of a group G is finite, then G is finite.*

Proof: Firstly, we have G is a torsion group, otherwise suppose $g \in G$ and $|g| = \infty$, then $\langle g^n \rangle$ will be different cyclic subgroups in G with infinitely many different n , a contradiction.

For any element g in G , it can generate a cyclic subgroup $\langle g \rangle$. Suppose G have n cyclic subgroups and $g_i (i = 1, \dots, n)$ be the generated elements of all the cyclic subgroups. $G = \cup_{i=1}^n \langle g_i \rangle$. So $|G| \leq \sum_{i=1}^n |g_i| < \infty$, that is to say G is finite.

3. Proof of the theorem

Proof of the theorem: We need only to prove the necessity of (2).

Case 1. G is abelian.

i) Suppose G is a torsion abelian group, we prove that G is a finite group.

Let $\pi(G)$ be the set of all primes dividing the orders of elements of G . For every prime p , let G_p be the set of all p -elements in G . Then G_p is a subgroup of G , that is, the p -component of G . We claim $|\pi(G)|$ is finite. If $|\pi(G)| = \infty$, then there must exist a prime $p \in \pi(G)$ such that G_p is a power subgroup for there are finitely many non-power subgroups. Hence there exists a positive integer m such that $G_p = G^m$, and then $G/G_p = G/G^m$, which means the exponent of G/G_p is finite. But $G/G_p \cong \prod_{q \in \pi(G) - \{p\}} G_q$. So the exponent of G/G_p can not be finite since $|\pi(G)|$ is infinite and then we get the contradiction.

If G_p is infinite for some p , then it contains infinitely many finite cyclic subgroups C_i and hence we may assume that C_i is a power subgroup in G and so G/C_i is of finite exponent, hence G is of finite exponent and G is a direct product of cyclic subgroups of finite order. In particular, G_p is of finite exponent and since $G_p \simeq G/N$ for some subgroup N , G_p has only finitely many non-power subgroups in G_p by Lemma 2. Since G_p has only finitely many power subgroups, i.e., $G_p, G_p^p, \dots, G_p^{p^l}$, where p^l is the power exponent of G_p , join with its finitely many non-power subgroups, G_p has only finitely many subgroups and so G_p is finite. Then for every $p \in \pi(G)$, G_p is finite. So G is finite.

ii) Suppose G is a non-torsion abelian group. Then G is an infinite abelian group with some elements of infinite order. We prove G is cyclic, which is contrary to $k > 0$ (k is the number of non-power subgroups of G).

Firstly, we prove that the subgroup T of G consisting of all elements of finite orders in G is finite. Evidently, no subgroup of T can be a power subgroup of G , otherwise G would be torsion and so, by i), G is finite, a contradiction. Hence, since G has only finitely many non-power subgroups, we have that T contains only finitely many subgroups. Thus T has only finitely many cyclic subgroups and T is finite by Lemma 4.

Now $T \neq G$ and $\overline{G} = G/T$ is an infinite torsion-free group. By Lemma 2, $\overline{G}$ has only finitely many non-power subgroups. We prove $\overline{G}$ is cyclic.

We claim that $\overline{G}/\overline{H}$ is finite for all non-trivial subgroup $\overline{H}$ of $\overline{G}$. In fact, let $\overline{H}$ be a nontrivial subgroup of $\overline{G}$. If $\overline{H}$ is a power subgroup of $\overline{G}$, then there exists a non-negative integer m such that $\overline{H} = \overline{G}^m$. Now $\overline{G}/\overline{H}$ is torsion, and it has only finitely many non-power subgroups by Lemma 2. Then $\overline{G}/\overline{H}$ is finite by i). On the other hand, suppose $\overline{H}$ is a non-power subgroup. Since $\overline{G}$ is torsion-free with finitely many non-power subgroups, there must exist $1 \neq \overline{H}_1 < \overline{H}$, with $\overline{H}_1$ being a power subgroup of $\overline{G}$. Then $\overline{G}/\overline{H}_1$ is finite and then $\overline{G}/\overline{H}$ is finite.

By Lemma 3, $\overline{G}$ is cyclic. Hence $G = T \times \langle z \rangle$ where z is of infinite order. It is obvious

that $G^m = T^m \times \langle z^m \rangle$, so if $s = |T| > 1$ then $T \times \langle z^{sn} \rangle$ is a non-power subgroup of G for each $n = 2, 3, \dots$. Thus $T = 1$ and so G is cyclic, a contradiction.

Case 2. G is non-abelian.

If G contains no non-power subgroups, then G is cyclic and every subgroup of a cyclic group is a power subgroup. So, suppose that G contains some non-power subgroups, and let $H_1, \dots, H_s$ be all of those. Since every conjugate of $H_i (i = 1, \dots, s)$ is also a non-power subgroup of G , so H_i has only finitely many conjugate subgroups. Then the normalizer N_i of H_i has a finite index in G and hence the subgroup $K_i = \cap_{g \in G} g^{-1} N_i g$ is normal in G and has a finite index in G . Let $K = \cap_{i=1}^s K_i$, then K is a normal subgroup of finite index in G and so it normalizes every H_i . Observe that every subgroup of K is normal in K . In fact, if $L \leq K$ and L is a power subgroup of G , then $L \trianglelefteq G$ and hence $L \trianglelefteq K$; if L is a non-power subgroup of G , then $L = H_i$ for some $i = 1, \dots, s$, and by construction of K , $H_i \trianglelefteq K$. Thus K is a Dedekind group. So K is either abelian or a direct product of an abelian group and the quaternion group of order 8. Hence the center $Z = Z(K)$ of K is of finite index in K and evidently $Z \triangleleft G$. Thus Z is an abelian normal subgroup of finite index, say m , in G . In particular, $x^m \in Z$ for every $x \in G$. Let l be the smallest natural number such that $x^l \in Z$ for every $x \in G$. Then $G^l \leq Z$.

Observe that G^l is a subgroup of finite index in Z . Otherwise, then Z/G^l is infinite and contains infinitely many proper non-trivial subgroups $R_i/G^l, i = 1, 2, \dots$. By assumption, there exists t such that R_t is a power subgroup of G , so $R_t = G^r$ for some $r \neq l$. It is easy to get $r < l$, but this contradicts the choice of l .

Thus the index of G^l in Z is finite, so we may replace Z by G^l and assume that Z is an abelian power subgroup of finite index in G , $Z = G^l$.

Observe that, by Lemma 1, every power subgroup of G , being contained in Z , is a power subgroup of Z , and every power subgroup of Z is a power subgroup of G . In particular, Z contains only finitely many non-power subgroups, and hence Z is finite or an infinite cyclic group. If Z is finite, then G is finite and so the result is proved. Thus we suppose Z is an infinite cyclic group, that is, $Z = \langle x \rangle$ with $|x| = \infty$. If $C_G(Z) \neq G$, then there exists $y \in G$ such that $x^y = y^{-1}xy \neq x \in \langle x \rangle$. Obviously $Z^y = Z$. Since x and x^{-1} are the only generators of $\langle x \rangle$, we have $x^y = x^{-1}$.

Let $Y_i = \langle x^{-i}yx^i \rangle, i = 1, 2, \dots$. We claim there are infinite many different Y_i . Suppose it is false. Let $Y_{i_1}, \dots, Y_{i_s}$ be all the different such type subgroups. Then there are infinite subgroups of this type equal to $Y_{i_t}, 1 \leq t \leq s$. Without any loss, we suppose $Y_{i_t} = Y_{m_1} = \dots = Y_{m_r} = \dots$. If $|y| = \infty$, by the generator properties of infinite cyclic group we have $y^{x^{m_1}} = (y^{x^{m_r}})^{\varepsilon_r}, r = 2, 3, \dots$, where $\varepsilon_r = 1$ or -1 . Hence there must exist r_0, r_1 such that

$\varepsilon_{r_0} = \varepsilon_{r_1}$ and $r_0 \neq r_1$. And then $y^{x^{m_{r_0}}} = y^{x^{m_{r_1}}}$. Suppose $m_{r_0} < m_{r_1}$ without any loss. Then we have $y = y^{x^{m_{r_1}-m_{r_0}}}$. But since $x^y = x^{-1}$, we get $y^{x^{m_{r_1}-m_{r_0}}} = x^{-2(m_{r_1}-m_{r_0})}y$. Therefore $y = y^{x^{m_{r_1}-m_{r_0}}} = x^{-2(m_{r_1}-m_{r_0})}y$, and then $x^{-2(m_{r_1}-m_{r_0})} = 1$. We get contradiction for $|x| = \infty$. On the other hand if $|y| = n$. Similarly there are infinite subgroups such that $Y_{m_1} = \cdots = Y_{m_r} = \cdots$. By the generator properties of cyclic group we have $y^{x^{m_1}} = (y^{x^{m_r}})^{\varepsilon_r}, r = 2, 3, \dots$, where $1 \leq \varepsilon_r \leq n$ is coprime to n . Hence there must exist r_0, r_1 such that $\varepsilon_{r_0} = \varepsilon_{r_1}$ and $r_0 \neq r_1$. And then $y^{x^{m_{r_0}}} = y^{x^{m_{r_1}}}$. We get a contradiction in the same way as before.

So there are infinite many different Y_i which are conjugate with each other. In particular, these infinite many Y_i are non-normal in G and non-power subgroups of G , which is contrary to the assumption. Thus $C_G(Z) = G$ and hence $Z \leq Z(G)$. So the center of G has a finite index in G . By Schur Theorem ([1] Theorem 10.1.3), the commutator subgroup $C = [G, G]$ is finite. If $C = 1$ then G is abelian contrary to the assumption. So let y be an element of prime order p in C . Obviously $Z \cap \langle y \rangle = 1$ and $\langle y \rangle \triangleleft \langle Z, y \rangle$ by $Z \leq Z(G)$, so $\langle Z, y \rangle = Z \times \langle y \rangle$ and all subgroups $Z^r \times \langle y \rangle$ are distinct for $r = 1, 2, \dots$. Thus there exists r such that $A = Z^r \times \langle y \rangle$ is a power subgroup of G . A contains, by Lemma 1, only a finitely many non-power subgroups of A and hence A is finite or cyclic. But this is not true. The theorem is proved.

Acknowledgement The authors would like to thank Prof. V.D. Mazurov for his help.

References

- [1] D.J.S. Robinson , An Course in the Theory of Groups, Springer-Verlag,Berlin 1982
- [2] F.Szász, On cyclic groups, Fund. Math., 43(1956), 238-240
- [3] Wei Zhou Lai ping, A characterization of non-cyclic groups, J. Southwest China Normal Univ. (Natural Sci.)(in Chinese), 2000, 25(3), 213-216.
- [4] Wei Zhou, An investigation on the least bound of the number of non-exponent subgroups of non-cyclic groups, ICM 2002, Abstracts of Short Comm. and Posters, Higher Education Press, Beijing, p.40